



The Series of Plates
Designed as a
Test for Colour-Deficiency

by

SHINOBU ISHIHARA M. D., Dr. Med. Sc.
Professor Emeritus of the University of Tokyo
Member of the Japan Academy

24 Plates Edition

KANEHARA TRADING INC.
TOKYO · JAPAN

Introduction

This series of plates is designed to provide a test which gives a quick and accurate assessment of colour vision deficiency of congenital origin. This is the commonest form of colour vision disturbances.

Most cases of congenital colour vision deficiency are characterized by a red-green deficiency which may be of two types; first, a protan type which may be absolute (protanopia) or partial (protanomalialia), and secondly, a deutan type which may be absolute (deutanopia) or partial (deutanomalialia).

In protanopia, the visible range of the spectrum is shorter at the red end compared with that of the normal, and that part of the spectrum which appears to the normal as blue-green, appears to those with protanopia as grey. The whole visible range of the spectrum in protanopia consists of two areas which are separated from each other by this grey part. Each area appears to those with protanopia as one system of colour with different brightness and saturation within each area, the colour in one area being different from that of the other. The red with a slight tinge of purple which is the complementary colour of blue-green appears also as grey.

In deutanopia, that part of the spectrum which appears to the normal as green, appears as grey, and the visible range of the spectrum is divided by this zone into two areas, each of which appears to be of one system of

colour. The visible range of the spectrum is not contracted, in contrast to protanopia. Purple-red which is the complementary colour of green appears also as grey.

In protanomalopia and deuteranomalopia, there is no part of the spectrum which appears as grey, but the part of spectrum which appears to those with protanopia as grey, appears to those with protanomalopia as a greyish indistinct colour, and likewise, the grey part of the spectrum seen by the person with deuteranopia appears to those with deuteranomalopia as an indistinct colour close to grey.

Consequently, one of the peculiarities of red-green deficiencies is that blue and yellow colours appear to be remarkably clear compared with red and green colours. The application of this peculiarity to the test for colour vision deficiencies is the distinguishing feature of this series.

In the congenital colour vision deficiencies, although very rare, there is total colour blindness which may be typical or atypical. The subject who suffers from typical total colour blindness shows a complete failure to discriminate any colour variations, usually with an associated impairment of central vision with photophobia and nystagmus. In the atypical total colour blindness, the colour sensitivity to red and green, as well as to yellow and blue is so low that only very clear colours can be perceived; but, except for the colour sensitivity, there is no abnormality in the visual functions. The plates in this book form an easy method of establishing the diagnosis on such cases and in distinguishing them from cases of red-green deficiencies.

Furthermore, a failure in the appreciation of blue and yellow may be termed tritanomalopia if partial, and tritanopia if absolute, but, even if such cases do exist, they are extremely rare. The plates in this book are not designed for the diagnosis of such cases.

How to use the test

The plates are designed to be appreciated correctly in a room which is lit adequately by daylight. The introduction of direct sunlight or the use of electric light may produce some discrepancy in the results because of an alteration in the appearance of shades of colour. When it is convenient only to use electric light, it should be adjusted as far as possible to resemble the effect of natural daylight. The plates are held 75 cm from the subject and tilted so that the plane of the paper is at right angle to the line of vision. The correct position of each plate is indicated by the number which is printed on the back of the plate. The numerals which are seen on plates 1—17 are stated, and each answer should be given without more than three seconds delay. If the subject is unable to read numerals, plates 18—24 are used and the winding lines between the two X's are traced with the brush. Each tracing should be completed within ten seconds.

It is not necessary in all cases to use the whole series of plates. Plates 16 and 17 may be omitted if the test is designed merely to separate the colour defectives from

those with normal colour appreciation. In a large scale examination the test may be simplified to an examination of six plates only; No. 1, one of Nos. 2, 3, one of Nos. 4, 5, 6, 7, one of Nos. 8, 9, one of Nos. 10, 11, 12, 13, and one of Nos. 14, 15.

It may be necessary to vary the order of the plates if it is suspected that there is a deliberate deception on the part of the subject.

Explanation of the plates

This series of plates is made up of the following 24 plates.

- No. 1. Both the normal and those with all sort of colour vision deficiencies read it as 12.
- Nos. 2, 3. The normal read them as 8 (No. 2) and 29 (No. 3). Those with red-green deficiencies read them as 3 (No. 2), and 70 (No. 3). Those with total colour blindness can not read any numeral.
- Nos. 4-7. The normal read them as 5 (No. 4), 3 (No. 5), 15 (No. 6) and 74 (No. 7). Those with red-green deficiencies read them as 2 (No. 4), 5 (No. 5), 17 (No. 6) and 21 (No. 7). Those with total colour blindness can not read any numeral.
- Nos. 8, 9. The normal read them as 6 (No. 8) and 45 (No. 9). The majority of those with colour vision deficiencies can not read them or read them incorrectly.

Nos. 10-13. The normal read them as 5 (No. 10), 7 (No. 11), 16 (No. 12) and 73 (No. 13). The majority of those with colour vision deficiencies can not read them or read them incorrectly.

Nos. 14, 15. The majority of those with red-green deficiencies read them as 5 (No. 14) and 45 (No. 15). The majority of the normal and those with total colour blindness can not read any numeral.

Nos. 16, 17. The normal read them as 26 (No. 16) and 42 (No. 17). In protanopia and strong protanomalopia only 6 (No. 16) and 2 (No. 17) are read, and in case of mild protanomalopia both numerals on each plate are read but the 6 (No. 16) and 2 (No. 17) are clearer than the other numerals. In deuteranopia and strong deuteranomalopia only 2 (No. 16) and 4 (No. 17) are read, and in case of mild deuteranomalopia both numerals on each plate are read but the 2 (No. 16) and 4 (No. 17) are clearer than the other numerals.

No. 18. In tracing the winding lines between the two X's, the normal trace along the purple and red lines. In protanopia and strong protanomalopia only the purple line is traced, and in case of mild protanomalopia both lines are traced but the purple line is easier to follow. In deuteranopia and strong deuteranomalopia only the red line is traced, and in case of mild deuteranomalopia both lines are traced but the red line is easier to follow.

- No. 19. In tracing the winding line between the two X's, the majority of those with red-green deficiencies trace along the line, but the majority of the normal and those with total colour blindness are unable to follow the line.
- No. 20. In tracing the winding line between the two X's, the normal trace the bluish-green line, but the majority of those with colour vision deficiencies are unable to follow the line or follow a line different from the normal one.
- No. 21. In tracing the winding line between the two X's, the normal trace the orange line, but the majority of those with colour vision deficiencies are unable to follow the line or follow a line different from the normal one.
- No. 22. In tracing the winding line between the two X's, the normal trace the line connecting the bluish-green and yellowish-green, those with red-green deficiencies trace the line connecting the bluish-green and purple, and those with total colour blindness can not trace any line.
- No. 23. In tracing the winding line between the two X's, the normal trace the line connecting the purple and orange, those with red-green deficiencies trace the line connecting the purple and bluish-green, and those with total colour blindness and weakness can not trace any line.
- No. 24. Both the normal and those with all sort of colour

vision deficiencies can trace the winding line between the two X's.

The typical readings of the normal person and the person with colour vision deficiencies on plates 1 to 25 are listed on the last page.

Analysis of the results

An assessment of the readings of plates 1 to 15 determines the normality or defectiveness of colour vision. If 13 or more plates are read normally, the colour vision is regarded as normal. If only 9 or less than 9 plates are read normally, the colour vision is regarded as deficient. However, in reference to plates 14 and 15, only those who read the numerals 5 and 45 and read them easier than those on plates 10 and 9 are recorded as abnormal readings.

It is rare to find a person whose recording of normal answers is between 14—16 plates. An assessment of such a case requires the use of other colour vision tests, including the anomaloscope.

In the assessment of colour appreciation by the short method involving 6 plates only as described on page 4, a normal recording of all plates is proof of normal colour vision. If there is a discrepancy in any of the recordings, the full series of plates should be used before diagnosing a red-green deficiency.

Care of the plates

It is important that the book of test plates should be kept closed, except during use, because undue exposure to sunlight causes a fading of the colour of the plates.

Plate	Normal Person	Person with Red-Green Deficiencies				Person with Total Colour Blindness
		Protan		Deutan		
		Strong	Mild	Strong	Mild	
1	12		12			12
2	8		3			X
3	29		70			X
4	5		2			X
5	3		5			X
6	15		17			X
7	74		21			X
8	6		X			X
9	45		X			X
10	5		X			X
11	7		X			X
12	16		X			X
13	73		X			X
14	X		5			X
15	X		45			X
		Protan		Deutan		
		Strong	Mild	Strong	Mild	
16	26	6	(2) 6	2	2 (6)	
17	42	2	(4) 2	4	4 (2)	

The mark X shows that the plate cannot be read. Blank space denotes that the reading is indefinite. The numerals in parenthesis show that they can be read but they are comparatively unclear.